

Wnt Genes

Review

Roel Nusse* and Harold E. Varmus†

*Howard Hughes Medical Institute
and Department of Developmental Biology
Stanford University School of Medicine
Stanford, California 94305

†Department of Microbiology and Immunology
and Department of Biochemistry and Biophysics
University of California, San Francisco
San Francisco, California 94143

Proto-Oncogenes and Development

During the past decade an avalanche of evidence has implicated proto-oncogenes in the control of cell proliferation. Products of proto-oncogenes include many of the secreted growth factors and cell surface receptors that mediate intercellular signaling events (Aaronson, 1991), the components of cytoplasmic pathways that interpret those signals (Cantley et al., 1991), and the nuclear factors that execute their commands through transcriptional regulation (Lewin, 1991). We are now witnessing a similar outpouring of support for the parallel notion that genes important in carcinogenesis are central to pattern formation in the embryo or differentiation in cell lineages. Preexisting or man-made mutations of mouse proto-oncogenes affect a variety of developmental processes (Forrester et al., 1992). Well-established growth factors and oncogenes can induce differentiation under appropriate conditions (e.g., Alema et al., 1985). The homologs of proto-oncogenes in flies, worms, and yeast are essential components of developmental mechanisms in those organisms (Hoffmann et al., 1992), and a few genes isolated as determinants of development, such as homeobox-encoding genes, have later been shown to have oncogenic potential (Perkins et al., 1990; Rabbitts, 1991).

Among the most striking connections between oncogenesis and development are provided by *Wnt* genes, the subject of this review. (The term *Wnt* is an amalgam of *wingless* [*wg*] and *int* [see below and Nusse et al., 1991].) The first *Wnt* gene was cloned from the mouse genome as a relatively obscure proto-oncogene ten years ago (Nusse and Varmus, 1982). But the numerous *Wnt* genes isolated from diverse species in the past few years have now attracted unusual attention because of the dramatic developmental phenomena attributed to them. Ectopic expression of *Wnt* genes induces axis duplication in frog embryos (McMahon and Moon, 1989), as well as mammary cancer in mice; *Wnt* gene deficiencies prevent normal development of mammalian brains (McMahon and Bradley, 1990; Thomas and Capecchi, 1990) and normal segmentation of insect embryos (Rijsewijk et al., 1987a).

Any effort to understand these remarkable effects must begin with the fact that *Wnt* genes encode secretory glycoproteins and apparently cause cells to proliferate, to differentiate, or perhaps simply to survive by signaling through autocrine and paracrine routes. In this sense, *Wnt* proteins are multipotent factors, capable of inducing different bio-

logical responses in different cellular contexts, a phenomenon also described for better-known secretory proteins such as the fibroblast growth factors (FGFs), the transforming growth factors β (TGF β), and nerve growth factor (NGF). Establishing a more detailed picture of the actions of *Wnt* genes, however, presents considerable challenges. The *Wnt* gene family is large, suggesting functional redundancy, the patterns of expression are complex, proteins encoded by *Wnt* genes have resisted isolation in biologically active form, and receptors for *Wnt* proteins, presumably essential components in cell–cell signaling, remain to be identified.

The *Wnt* Gene Family

With the benefit of hindsight, we now recognize that phenomena studied for several decades are the consequences of *Wnt* gene mutations. Viral insertion mutations regularly promote mammary tumors in laboratory mice (Bittner, 1936; Korteweg, 1936), a spontaneous frameshift mutation of mice (*swaying*) impairs cerebellar structure and function (Lane, 1967; Thomas et al., 1991), and *wg* mutations in *Drosophila* can transform a wing to a notum or disrupt segment polarity (Sharma, 1973; Nüsslein-Volhard and Wieschaus, 1980).

Wnt genes were initially cloned as candidate proto-oncogenes (mouse *Wnt-1* and *Wnt-3* [Nusse and Varmus, 1982; Roelink et al., 1990]), as a gene near the cystic fibrosis locus (human *Wnt-2*; Wainwright et al., 1988), and as human, *Drosophila*, and *Xenopus* homologs of mouse *Wnt-1* (Van't Veer et al., 1984; Rijsewijk et al., 1987a; Noordermeer et al., 1989). Once these few *Wnt* genes were in hand to identify conserved sequences for primers, the polymerase chain reaction (PCR) produced an outpouring of *Wnt* gene family members. Up to ten or more *Wnt* coding sequences have been isolated from the widely studied species listed in Table 1, as well as from zebrafish (Molven et al., 1991), several other vertebrates (including birds, reptiles, and jawless fishes), sea urchins, starfish (Sidow, 1992), *Axolotl* (Busse et al., 1990), and leech (Kostriken and Weisblat, 1992). It is doubtful whether the full *Wnt* repertoire is in hand for any of these species. Even in the mouse, which has yielded the largest number of published sequences, most were recovered from cDNA libraries of midgestational embryonic RNA with a single set of PCR primers (Gavin et al., 1990). Other libraries and other primers may well produce more genes.

Each of the fully sequenced *Wnt* open reading frames encodes a protein that appears destined for secretion, harbors one or more sites for N-linked glycosylation, and conserves up to 23 or 24 cysteines in nearly parallel positions (Figure 1). Most of the deduced proteins are about 350 to 380 aa in length, with over 100 conserved residues fairly evenly distributed across the entire sequence. Some *Wnt* proteins have additional internal, amino-terminal, or carboxy-terminal domains; for example, the *wg* protein has 85 aa inserted just before the region encoded by its final exon, the DWnt-3 protein has a 155 aa insertion and

Table 1. *Wnt* Genes Identified in Commonly Studied Organisms, with Chromosomal Assignments


Species	Genes	Chromosome Assignments	References
<i>Homo sapiens</i>	<i>Wnt-1</i>	12q13	Van't Veer et al., 1984; van Ooyen et al., 1985
	<i>Wnt-2</i>	7q3.1	Wainwright et al., 1988
	<i>Wnt-3</i>	17q21	R. N., unpublished data
<i>Mus musculus</i>	<i>Wnt-1</i>	15	Nusse and Varmus, 1982; Nusse et al., 1984; van Ooyen and Nusse, 1984; Fung et al., 1985
	<i>Wnt-2</i>	6	McMahon and McMahon, 1989
	<i>Wnt-3</i>	11	Roelink et al., 1990
	<i>Wnt-3A</i>	11	Roelink and Nusse, 1991
	<i>Wnt-4</i>	4	Gavin et al., 1990
	<i>Wnt-5A</i>	14	Gavin et al., 1990
	<i>Wnt-5B</i>	6	Gavin et al., 1990
	<i>Wnt-6</i>	1	Gavin et al., 1990
	<i>Wnt-7A</i>	6	Gavin et al., 1990
	<i>Wnt-7B</i>	15	Gavin et al., 1990
<i>Xenopus laevis</i>	<i>X-Wnt-1</i>	?	Noordermeer et al., 1989
	<i>X-Wnt-3</i>	?	Christian et al., 1991a, 1991b
	<i>X-Wnt-4</i>	?	Christian et al., 1991a, 1991b
	<i>X-Wnt-5A</i>	?	Christian et al., 1991a, 1991b
	<i>X-Wnt-8</i>	?	Christian et al., 1991a, 1991b
<i>Drosophila melanogaster</i>	wg	28A	Rijsewijk et al., 1987a
	D-Wnt-2	45E	Russell et al., 1992
	D-Wnt-3	17B	Russell et al., 1992
<i>C. elegans</i>	Ce-Wnt-1	II	Kamb et al., 1989
	Ce-Wnt-2	IV	R. Waterston, personal communication

a long amino-terminal extension, and the mouse *Wnt-2* and *Xenopus Wnt-8* proteins have short carboxy-terminal extensions. In most cases, *Wnt* proteins display 30%–60% amino acid identity when two proteins within a single species or two from widely divergent species are compared. In fact, *Wnt* genes are sufficiently well conserved to permit recognition of the homologs of individual mouse genes in different orders of vertebrates (e.g., Christian et al., 1991a) and even in insects or echinoderms (Sidow, 1992). For example, as pointed out by Gavin et al. (1990), mouse *Wnt-1* protein and its orthologs in *Xenopus*, zebrafish, and *Drosophila* share one cysteine residue in the amino-terminal region that is not present in other family members; conversely, other *Wnt* proteins contain two nearby cysteine residues not found in *Wnt-1* proteins (Figure 1).

Based upon such arguments it appears that at least four

and perhaps as many as seven *Wnt* genes must have existed before the divergence of arthropods and chordates (Sidow, 1992). In addition, amino acid substitutions in *Wnt* protein sequences have accumulated at an especially slow rate during vertebrate evolution (Sidow, 1992), consistent with the central inductive roles proposed for *Wnt* proteins during embryogenesis. Even though some pairs of proteins (e.g., mouse *Wnt-3* and -3A, *Wnt-5A* and -5B, or *Wnt-7A* and -7B) are very highly related, the appearance of these pairs in cartilaginous fish implies that the duplications producing them must still have occurred at least 400 million years ago. As might be expected under these circumstances, there are very few differences between the *Wnt* proteins of mouse and human that have been compared (van Ooyen et al., 1985; McMahon and McMahon, 1989).

Limited information is available about the organization

Figure 1. Schematic Representation of Conserved Amino Acid Residues in *Wnt* Proteins

Conserved cysteine residues are represented by continuous bars; a cysteine unique for *Wnt-1* and *wg* is marked by a plus sign; and two cysteines found in all other *Wnt* proteins by asterisks. Other absolutely conserved residues are shown as wavy bars, and residues conserved in at least 75% of *Wnt* proteins by striped bars. A signal sequence (ss) is present at the amino terminus. Inserts in *Wg* and *DWnt-3* proteins are shown below. The figure is based on an alignment provided by A. Sidow.

Table 2. Expression Patterns of *Wnt* Genes in the Mouse

Gene	Expression in Embryos	Expression in Adults	Expression in Mammary Gland
<i>Wnt-1</i>	Brain; ring around midbrain-hindbrain junction; tip headfolds; dorsal wall diencephalon; roofplate spinal cord	Testes (postmeiotic spermatids)	Not detectable
<i>Wnt-2</i>	Developing allantois; pericardium heart; ventral-lateral mesoderm	Lung; brain; heart; placenta	Virgins
<i>Wnt-3</i>	D2 neuromer diencephalon (dorsal thalamus); ventral horns spinal cord; developing cerebellum; ectoderm overlying mandible, genital tubercle, and limbs	Thalamus; Purkinje cells in cerebellum; pons; Shaft hair root in skin	Not detectable
<i>Wnt-3A</i>	Ectoderm and mesoderm primitive streak posterior hindbrain; anterior midbrain; roofplate spinal cord; archicortex of the telencephalon	Lung	Not detectable
<i>Wnt-4</i>	Embryo; fetus (sites not determined)	Brain; lung	Virgins; early pregnancy
<i>Wnt-5A</i>	Face; brain (ventral area midbrain) and spinal cord; gradient in developing limbs	Heart; lung	Early pregnancy
<i>Wnt-5B</i>	Embryo; fetus, not restricted	Heart; liver; brain; lung; testes; kidney	Midpregnancy
<i>Wnt-6</i>	Embryo; fetus (sites not determined)	Testes	Midpregnancy, early lactation
<i>Wnt-7A</i>	Embryo; fetus (sites not determined)	Brain; lung	Not detectable
<i>Wnt-7B</i>	Embryo; fetus (sites not determined)	Brain	Virgins; early pregnancy

of *Wnt* genes and the structural determinants of their expression. Of the several genes that have been sequenced, most have three introns in corresponding positions; however, a *Caenorhabditis elegans* *Wnt* gene has a radically different distribution of exons and introns (G. Shackleford, L. Shiue, H. E. V., unpublished data), and a *Drosophila* gene, *DWnt-3*, has no introns at all (Russell et al., 1992). There is fragmentary evidence for alternative modes of expression: multiple sites for initiation of transcription (Nusse et al., 1990) and for polyadenylation (Gavin et al., 1990) and at least one example of alternative splicing to produce two slightly different proteins from mouse *Wnt-4* (Gavin et al., 1990). However, the important determinants of the complex transcriptional programs described below (Tables 2 and 3) have not been defined.

Wnt Proteins and Their Signaling Mechanism

The predicted primary protein products of *Wnt* genes display many of the characteristics of secreted growth factors: a hydrophobic signal peptide, followed by a recognition site for signal peptidase, the absence of additional transmembrane domains, abundant and strongly conserved cysteine residues with potential for disulfide cross-linking, and prospective sites for N-linked glycosylation (Figure 1).

Despite these strong clues to the nature of *Wnt* proteins, it has been difficult to document that they are secreted and thus far impossible to prepare them in a useful cell-free form. Most of the available information about *Wnt* proteins has been obtained from the study of the mouse *Wnt-1*, human *Wnt-2*, and the *Drosophila* *wg* genes, largely because antisera have been prepared to detect their products. Since tissues that express endogenous *Wnt-1* are not readily cultured and since nearly all cultured cell lines (excepting embryonal carcinoma lines such as P19 [St. Arnaud et al., 1989; Schuurin et al., 1989]) fail to express the gene, most work on *Wnt-1* proteins has been performed with cells programmed to express exogenous cDNA under the control of a heterologous promoter.

In these ectopic settings, a large portion of *Wnt-1* and *Wnt-2* protein remains associated with internal membranous components of cells, in a series of glycosylated forms, ranging from 36 to 42 kd in size (Brown et al., 1987; Papkoff et al., 1987; Blasband et al., 1992). It is likely that most of the *Wnt* protein is in the endoplasmic reticulum (ER) and cis-Golgi apparatus; intracellular *Wnt-1* proteins are associated with the 78 kd, ER-based chaperonin called BiP (Kitajewski et al., 1992), and the carbohydrate modifications are immature (susceptible to endoglycosidase H [Papkoff, 1989]).

Table 3. *Wnt* Genes in Xenopus

Gene	Expression	Duplication Axis	Effects on Gap Junctions
<i>XWnt-1</i>	Neurula	Anterior	Ventral opening
<i>XWnt-3A</i>	Neurula	Partial anterior	
<i>XWnt-4</i>	Neurula (dorsal in brain) and floor plate spinal cord		
<i>XWnt-5A</i>	Oocyte/tadpole (maternal)	Posterior	No effect
<i>XWnt-8</i>	Midblastula; band in marginal zone (excluding organizer)	Anterior	Ventral opening




Figure 2. Direct and Paracrine Cell Culture Assays for *Wnt* Genes

In a direct assay for *Wnt* genes, nonresponsive cells (circles) or mammary epithelial cells (squares) that respond to *Wnt* genes by morphological and proliferative changes (oblongs) produce *Wnt* proteins (as indicated by stippling) after introduction of a *Wnt* expression vector. In one version of a paracrine assay, nonexpressing or *Wnt*-expressing colonies of nonresponsive cells are surrounded by untreated responders. Photographs at the bottom show, on the left, C57MG cells before and after introduction of a retrovirus vector carrying *Wnt-1*, and, on the right at lower magnification, nonexpressing C57MG cells surrounding colonies of rat-1 cells that are or are not infected with the virus vector bearing *Wnt-1*. The photographs are from Jue et al. (1992) and J. Kitajewski and H. E. V.

Under normal conditions, *Wnt* proteins have not been detected by either immunological or biological assays in culture medium from cells expressing exogenous *Wnt* genes. But immunoreactive forms can be recovered by treating the cultures with the polyanion, suramin (Papkoff, 1989; Papkoff and Shryver, 1990; Blasband et al., 1992), by solubilizing the extracellular matrix after removing the cells, or by growing the cultures in the presence of soluble heparin, to which *Wnt* proteins bind at physiological salt concentrations (Bradley and Brown, 1990). *Wnt* proteins identified in these ways are apparently secreted, although at different efficiencies in different cell lines, and their mobilities in gel electrophoresis suggest varying degrees of further modification. Based upon antibody binding and iodination experiments with intact cells (Papkoff and Shryver, 1990), *Wnt* proteins appear to be associated with the cell surface, as well as with the extracellular matrix. However, it is not known whether these associations are specific, whether certain forms of the protein (e.g., oligomers or heterologous complexes) are biologically active, or whether components of the extracellular matrix, such as glycosaminoglycans, can act as cofactors to promote the response to *Wnt* proteins, in the manner recently demonstrated for FGFs (Klagsbrun and Baird, 1991).

Despite the failures to obtain *Wnt-1* protein in a biologically active, cell-free form, the secreted protein appears to have biological activity, as measured in cell transformation assays that depend upon paracrine effects (Figure 2; Jue et al., 1992). In these tests, transforming activity can be supplied by any of several cell lines that expresses an introduced *Wnt-1* gene without detectably responding to it (e.g., mouse 3T3, rat-1, quail sarcoma, or HeLa cells). When *Wnt*-responsive C57MG cells, derived from a normal mouse mammary gland, are mixed with or surround the donor cells, they undergo morphological changes indistinguishable from those produced when a *Wnt-1* vector is expressed directly in C57MG cells (see below and Figure 2).

Although available antibodies against *Wnt-1* proteins do not block the paracrine effect, it seems very likely to be attributable to *Wnt-1* proteins: many cells can serve as

donors, yet the only responders are the mammary cells that are also susceptible to direct transformation by *Wnt-1* and related genes. In addition, the effect is blocked by heparin (Jue et al., 1992), and it is not observed with a *Wnt-1* mutant whose product lacks the signal peptide (Mason et al., 1992). This mutant also fails to transform C57MG cells directly (Mason et al., 1992) or to cause axis duplication in *Xenopus* embryos (McMahon and Moon, 1989); thus, entry into the secretory pathway appears to be essential for biological activity of *Wnt* proteins. In *Drosophila*, where genetic evidence exists for the paracrine activity of *wg* (see below), immunostaining of embryos has been used to show that *Wg* protein arrives at the surface of cells containing *wg* RNA and can be found in adjacent cells (van den Heuvel et al., 1989).

The implied importance of the conserved cysteine residues in *Wnt* proteins has been further documented by the study of *Wnt-1* mutants. When the conserved cysteine codon at the penultimate position in mouse *Wnt-1* is changed to a tryptophan codon (Cys369Trp), the gene loses its ability to transform C57MG cells directly or by a paracrine mechanism (Mason et al., 1992) and to induce axis duplication in *Xenopus* embryos (McMahon and Moon, 1989). Mutation of other cysteine residues also impairs transforming activity; in one case (Cys151Ser) transformation of C57MG cells is temperature sensitive (Mason et al., 1992). Curiously, the same change at a nonhomologous cysteine residue in *wg* protein (Cys107Ser) is responsible for a spontaneous temperature-sensitive mutant of *wg* (M. van den Heuvel, C. Harryman, and R. N., unpublished data). With one exception, glycosylation sites, in contrast to cysteine residues, are not well conserved, and none of the four sites in *Wnt-1* protein is required for transforming activity; indeed, a mutant protein lacking all four sites is still transformation competent (Mason et al., 1992). However, one glycosylation site mutant (Asn359Gln) is also temperature sensitive for transformation.

***Wnt* Genes in Mammary Carcinogenesis**

Wnt genes are sources of differentiation-inducing signals during normal developmental events, but they also have

the potential to promote carcinogenesis through local effects on cell proliferation, particularly in the mammary gland. Indeed, the first *Wnt* gene to be molecularly cloned, mouse *Wnt-1*, was isolated during a search for cellular genes insertionally activated by proviral DNA in mammary carcinomas induced by the mouse mammary tumor virus (MMTV) (Nusse and Varmus, 1982). (Until recently [Nusse et al., 1991], it was called *int-1*, for MMTV integration site.) Although transcription of the *c-myc* proto-oncogene, the progenitor of the retroviral oncogene, *v-myc*, had earlier been shown to be stimulated by adjacent avian leukosis virus DNA in chicken B-cell lymphomas (Hayward et al., 1981), *Wnt-1* was the first of many proto-oncogenes to be discovered originally as a target for proviral insertion mutations (van Lohuizen and Berns, 1990). The proviral tagging method has subsequently identified at least four additional proto-oncogenes that are transcriptionally activated by MMTV proviruses in mouse mammary carcinomas: two members of the FGF gene family, *int-2* (also known as *FGF-3*; Dickson et al., 1984; Dickson and Peters, 1987) and *hst-1* (also known as *K-fgf* or *fgf-4*; Peters et al., 1989), another member of the *Wnt* gene family (*Wnt-3*; Roelink et al., 1990), and *int-3*, a gene encoding a presumptive transmembrane receptor with regions homologous to the products of *Notch*, *lin-12*, and *Tan-1* (Gallahan and Callahan, 1987; Robbins et al., 1992). In a provocatively high percentage of tumors, insertions have occurred in two of these genes, usually *Wnt-1* and *int-2* (Peters et al., 1986; Mester et al., 1987), suggesting cooperative effects in carcinogenesis (see below). In addition, a third *Wnt* gene, *Wnt-2*, has been implicated in mammary tumorigenesis because it is amplified and overexpressed in subclones of two transplanted, virus-induced tumors (Roelink et al., 1992).

Notably, none of the MMTV-activated genes has been encountered as a naturally occurring retroviral oncogene; only *hst* has been implicated as a proto-oncogene in other contexts (Taira et al., 1987), and all five appear to affect extracellular signaling events—four by production of secretory proteins and one by production of a presumptive cell surface receptor. These observations are surprising, since genes such as *c-myc* and *c-Ha-ras*—which have been repeatedly transduced by retroviruses, are involved in many kinds of tumors, and encode intracellular regulators—do have the potential to induce mammary carcinomas when engineered to form a transgene under the control of an MMTV long terminal repeat (Stewart et al., 1984; reviewed by Adams and Cory, 1991).

Initially, a causative role for *Wnt* genes in mammary carcinogenesis was based upon three circumstantial arguments: the repeated occurrence of proviral insertion mutations, very commonly in the *Wnt-1* locus, occasionally in *Wnt-3*; the consequent activation of expression of these genes, which are normally silent in the mammary gland, most often by provision of a viral enhancer that acts upon the normal *Wnt* promoter (Nusse et al., 1984, 1990; Roelink et al., 1990; Table 2); and the retention of a complete open reading frame, even when the insertions occur within exons (van Ooyen and Nusse, 1984; Roelink et al., 1990). Since *Wnt* loci are not known to be favored sites for integra-

tion, it is presumed that the transcription-activating insertions in the *Wnt-1* and *Wnt-3* loci confer a strong growth advantage upon individual cells, accounting for the clonal character of the tumors (Cohen et al., 1979).

These indirect arguments for the oncogenic effects of *Wnt* insertion mutations have been made more persuasive by gene transfer experiments in cell culture and animals. Although the mouse *Wnt-1* gene has no evident phenotypic effects upon primary embryo cells or most established cell lines, it can induce morphological changes and enhanced growth properties when expressed in at least two mouse mammary epithelial cell lines, C57MG (Brown et al., 1986) and RAC (Rijsewijk et al., 1987b). Similar findings have been obtained with human *Wnt-2* (Blasband et al., 1992), mouse *Wnt-3A* (N. Parkin and H. E. V., unpublished data), and *wg* (A. M. C. Brown, personal communication). (Indeed, only one other type of cultured cell, the rat pheochromocytoma line, PC12, is known to respond to *Wnt* genes [G. Shackleford and H. E. V., unpublished data].) The conversion of cuboidal, mammary epithelial cells to elongated, highly refractile cells that continue to replicate in a dense culture provides a useful assay for *Wnt* genes (Figure 2). The phenomena induced in mammary cell lines, however, do not necessarily simulate the changes anticipated for a mammary oncogene. Transformation of one of the lines (C57MG) is not accompanied by an ability to form tumors in animals (Brown et al., 1986), and cells taken directly from naturally occurring mouse mammary tumors more closely resemble the original C57MG and RAC cell lines than the *Wnt*-transformed cells (e.g., Sonnenberg et al., 1987).

Transgenic mice have provided conclusive evidence for the oncogenic potential of *Wnt-1*. Mice transmitting a *Wnt-1* transgene that mimics activated alleles in virus-induced tumors exhibit extensive hormone-independent hyperplasia of mammary epithelium, so that glands in virgin female and male animals resemble those of pregnant animals (Tsukamoto et al., 1988). By 1 year of age, mammary carcinomas indistinguishable from virus-induced tumors have appeared in most females, in one or two of the ten glands, and in about 15% of males. Occasional salivary gland carcinomas are also observed.

As in several other transgenic models for oncogenesis (Adams and Cory, 1991), the stochastic appearance of mammary carcinomas in *Wnt-1* transgenic animals suggests that additional events, presumably mutations affecting proto-oncogenes or tumor suppressor genes, are required for full oncogenic conversion of cells proliferating under the influence of the *Wnt-1* transgene. Genes capable of collaborating with an activated *Wnt-1* gene during mammary tumorigenesis have been sought by crossing lines of transgenic mice and by infecting transgenic mice with MMTV. Both approaches support earlier indications from studies of nontransgenic animals that the *Wnt-1* and *int-2* genes can act cooperatively to promote tumors. When *Wnt-1* and *int-2* transgenic animals are mated to produce bitransgenic animals, mammary tumors appear earlier and, in males, at a much higher frequency than in either type of parental animal (Kwan et al., 1992). Infection with MMTV also accelerates the onset of tumors in virgin

and breeding female mice carrying the *Wnt-1* transgene. Many of the tumors are derived from infected cells, and they exhibit proviral insertion mutations of *int-2* (most frequently), *hst-1*, or (occasionally) both genes (G. Shackelford, C. MacArthur, H. Kwan, and H. E. V., unpublished data). Other proviruses may have caused contributory mutations at additional, and perhaps novel, loci that have yet to be characterized. The frequent cooperation of *Wnt* and *FGF* gene family members in MMTV-induced mammary carcinogenesis is especially intriguing. *Wnt-1* and *int-2* proteins, although unrelated by sequence, are secretory glycoproteins with surprisingly similar biochemical properties (Kiefer et al., 1991), and *Wnt* and *FGF* proteins can also collaborate during the induction of *Xenopus* mesoderm in vitro (Christian et al., 1992; see below).

The pathogenesis of mammary tumors under the influence of an activated *Wnt* gene remains poorly understood. The appearance of mammary hyperplasia in *Wnt-1* transgenic mice indicates that *Wnt-1* protein can act as a growth stimulus in early phases of the carcinogenic process, but roles for *Wnt* genes in later stages—malignant conversion, escape from hormonal (e.g., pregnancy) dependence, or acquisition of metastatic potential—have not been defined. The *Wnt* genes most convincingly implicated in mammary tumorigenesis (*Wnt-1* and *Wnt-3*) are not normally expressed at detectable levels in the mammary gland; hence, receptors for their products might not be expected to be present on mammary cells. However, several other members of the *Wnt* gene family are expressed during maturation of the mammary gland (see below and Table 2). Thus, it is likely that the surfaces of mammary cells are equipped with either a general receptor for *Wnt* proteins or receptors for the normally produced *Wnt* proteins that can be usurped by products of the activated genes. It has not been established whether those *Wnt* genes that are normally active in the mammary gland can act as oncogenes if expressed at yet higher levels. Nor has an alternative proposal been formally excluded: that oncogenic *Wnt* proteins interfere with growth-restraining signals supplied by *Wnt* proteins normally found in the mammary gland.

Local effects upon cell growth by *Wnt* and other secretory proteins may occur by either an autocrine or a paracrine mechanism, but no evidence decisively distinguishes between these mechanisms in mammary tumorigenesis. Because mouse mammary tumors are composed mostly of clonal populations of cells bearing an insertionally activated *Wnt* or *FGF* locus, or both, the autocrine mode appears to be preferred. Nevertheless, in a few tumors with multiple mutations, transplantation has ultimately separated tumor cells with different mutations (Mester et al., 1987; Roelink et al., 1992), implying that the original tumors were oligoclonal and perhaps sustained in part by paracrine mechanisms.

Tumors induced by secretory factors, such as *Wnt* proteins or *FGFs*, may be useful models for human cancers in which growth factors or their receptors have been implicated in pathogenesis. To date, however, mutations of *Wnt* genes have not been reported in tumors other than mammary or salivary gland carcinomas in MMTV-infected

or transgenic mice. In particular, such mutations have not been observed in human breast cancer (Van de Vijver et al., 1989; Van de Vijver and Nusse, 1991), although the number of *Wnt* genes studied and the extent of the surveys have been limited. Two other targets for MMTV insertion mutations, *int-2* and *hst*, are sometimes amplified in human cancer, including breast cancer (e.g., Ali et al., 1989), but the amplified *FGF* genes are not expressed, and co-amplified genes (such as *bcl-1/PRAD1* [Lammie et al., 1991]) are also candidate oncogenes.

Expression of the *Wnt* Genes in the Mouse and the *Wnt-1* Phenotype

The highly restricted pattern of expression of *Wnt-1* in adult and embryonic mice (Jakobovits et al., 1986; Shackelford and Varmus, 1987; Wilkinson et al., 1987) was the first indication that *Wnt* genes might regulate major events in mammalian development. Since then, many *Wnt* family members have been shown to be expressed in equally provocative patterns during mouse embryogenesis (Gavin et al., 1990; McMahon and McMahon, 1989; Roelink and Nusse, 1991; Table 2). Although the lack of suitable antisera has prevented definition of *Wnt* protein distribution within tissues, the sharp boundaries observed by in situ hybridization to *Wnt* RNA suggest that the genes are involved in the establishment of specific cell fates in a regional manner.

This proposal has received its most resounding support from the phenotype of *Wnt-1* mutants in mice. Presumptive null mutations have been obtained in two ways: *Wnt-1^{neo}* alleles result from homologous targeting events that insert *neo* into the second exon of *Wnt-1* in embryonic stem cells (Thomas and Capecchi, 1990; McMahon and Bradley, 1990), whereas *Wnt-1^{sw}* is a naturally occurring single nucleotide deletion mutation that prematurely terminates translation in the middle of the *Wnt-1* open reading frame in *swaying* mice (Lane, 1967; Thomas et al., 1991). Mice homozygous for either *Wnt-1^{neo}* or *Wnt-1^{sw}* and double heterozygotes display a range of phenotypes. At one extreme, animals die around the time of birth, lacking the entire cerebellum and a significant portion of the midbrain (McMahon and Bradley, 1990; McMahon et al., 1992). The loss of these structures can be detected in embryos as early as day 9.5, although the remaining parts of the central nervous system (CNS), at that age and later, are remarkably intact. At the other extreme, some homozygous mutant animals survive into adulthood, suffering from ataxia and lacking the anterior half of the cerebellum (Thomas and Capecchi, 1990; Thomas et al., 1991). Thus, the region affected in *Wnt-1* mutants minimally includes the anterior half of the cerebellum, but often extends significantly into the midbrain or the posterior cerebellum. The explanation for the variable penetrance of the *Wnt-1* mutations is unknown, but may be due to several factors, including the genetic background.

The *Wnt-1* expression pattern in the developing CNS between days 8 and 14 of embryogenesis partially coincides with the regions affected in *Wnt-1* mutants (Figure 3 and Table 2; Wilkinson et al., 1987; McMahon et al., 1992). Around day E9.5, *Wnt-1* RNA is detected in a circle




Figure 3. Expression of *Wnt-1* in Mouse and Zebrafish Embryos and the *Wnt-1* Phenotype

The expression pattern of *Wnt-1* in the brain of an 8.5 day embryo (stippled in [A]) is compared with the area deleted in a newborn *Wnt-1* mutant (stippled area in [B]). The superior and inferior colliculi (sc, ic), derived from the midbrain (mes), and the cerebellum (cer), derived from the anterior hindbrain (metencephalon [met]), fail to develop in the example shown; but in some embryos, the midbrain and posterior half of the cerebellum remain intact, as discussed in the text. The forebrain (fb) and myelencephalon (my) seem to be unaffected. (C) and (D) show the pattern of expression (stippled area) of *Wnt-1* in the CNS of a 22 hr zebrafish embryo. Dorsal view (C); side view (D). Expression is seen in dorsal structures of the mesencephalon (mes), the hindbrain (h), and spinal cord (s), and in a ring anterior to the midbrain-hindbrain junction. As in the mouse, *Wnt-1* is not expressed in the anterior hindbrain and the forebrain (fb). (Reproduced with permission from Nature 347, p. 335, 1990, and from EMBO J. 10, p. 805, 1991.)

of cells in the most posterior portion of the midbrain, just anterior to the junction with the hindbrain (Wilkinson et al., 1987), in the region that contributes to the anterior half of the cerebellum (Hallonet et al., 1990). In addition, the gene is expressed at early times in the region that probably becomes the midbrain (or tectum) in the mature animal. A strikingly similar pattern of *Wnt-1* RNA in the zebrafish embryo, including the ring of cells in the midbrain, under-

scores the functional relevance of these expression sites (Molven et al., 1991; Figure 3). *Wnt-1* is not expressed, however, in the anterior hindbrain, the region from which the posterior half of the cerebellum is derived (Hallonet et al., 1990). It has been postulated that the absence of the entire cerebellum is an indirect effect of *Wnt-1*, mediated through the loss of expression of the mouse *engrailed* (*en*) genes (McMahon et al., 1992). In this scheme, *Wnt-1* would normally maintain *en* expression through an inductive event in adjacent tissue, in much the same way as found in *Drosophila* embryos (see below). Expression of *en* would then be essential for the correct outgrowth of the posterior half of the cerebellum.

Wnt-1 mutants appear to have a normal spinal cord and hindbrain, even though *Wnt-1* is normally expressed in the caudal part of the hindbrain and in the dorsal midline of the spinal cord from the earliest times that those structures are recognizable (Figure 3; Wilkinson et al., 1987; Molven et al., 1991; McMahon et al., 1992). The absence of a mutant phenotype at other *Wnt-1* expression sites in the CNS is probably due to complementing genes, particularly other *Wnt* genes. The *Wnt-3A* gene, which is expressed in the caudal hindbrain and the roofplate in the spinal cord, but not the posterior midbrain, is a leading candidate to complement the *Wnt-1* deficiency (Roelink and Nusse, 1991; McMahon et al., 1992).

In adult mice, *Wnt-1* is normally expressed only in the male germline, in the round spermatid stage of spermatogenesis (Shackleford and Varmus, 1987). Since at least some *swaying* males are fertile (P. Neumann, personal communication), *Wnt-1* is not absolutely required for male gametogenesis.

The sites of expression of most additional members of the *Wnt* family are still incompletely catalogued. Transcripts of many *Wnt* genes have been found in different, spatially restricted patterns in embryos, in most organs of adult animals, with the possible exception of lymphoid tissues, and in cells derived from all three germ layers (see Table 2; Gavin et al., 1990; McMahon and McMahon, 1989; Roelink and Nusse, 1991). Frequent expression of other *Wnt* genes during development of the CNS suggests functions similar to those provided by *Wnt-1*, at other times and locations (Table 2). *Wnt* genes may also be involved in normal development of the limbs. For instance, *Wnt-3* is expressed in the ectoderm overlying the limb buds (Roelink and Nusse, 1991), and *Wnt-5A* is expressed initially in the ectoderm and later in a gradient in the limb mesenchyme, with the highest levels at the distal end (Gavin et al., 1990). These patterns suggest a role in directing the growth and polarity of the limbs, possibly in conjunction with similar gradients of homeobox gene products (Dollé et al., 1989; Oliver et al., 1988). Pairs of *Wnt* genes highly related to each other (3 and 3A; 5A and 5B; 7A and 7B) usually have distinct, nonoverlapping expression sites (Gavin et al., 1990); for example, in contrast to *Wnt-5A*, *Wnt-5B* shows no gradients in limb buds, implying that very similar proteins act at different sites, perhaps through a common biochemical mechanism.

Although the role of *Wnt* genes in postnatal differentiation has yet to be carefully studied, special attention has




Figure 4. The Role of *Wnt* Genes in Early Frog Development

The upper half shows a current model of mesoderm induction in *Xenopus* embryos. A signal from the vegetal half (V1) induces mesoderm in the overlying animal cap of the blastocyst. At the dorsal side of the embryo, a special signal (V2) from the Nieuwkoop center induces a Spemann organizer (O) in the mesoderm. This organizer determines the axis of the embryo. The lower half shows that the injection of any of several *Wnt* RNAs results in the formation of a second axis through the generation of a second Spemann organizer. This could result from the injected cells acting as a second organizer (upper arrow; Sokol et al., 1991) or as a second Nieuwkoop center, which in turn induces a second organizer (lower pathway). In the latter case, the *Wnt*-injected cells would not be present in the induced second axis, consistent with the findings of Smith and Harland (1991).

been directed to the normal mammary gland, in view of the oncogenic effects of *Wnt* genes discussed earlier. At least six *Wnt* genes are expressed in the mammary gland, in a variety of temporal patterns, in virgin, pregnant, and lactating animals; none of these seems to be frequently activated in mouse mammary cancer (Gavin and McMahon, 1992; J. Rosen, personal communication; Table 2). Although difficulties with in situ hybridization techniques have impeded description of the spatial patterns, it seems likely that *Wnt* genes are important regulators of development in the mammary gland—and perhaps in other adult organs as well.

Inductive Properties of *Wnt* Genes in *Xenopus* Embryos

The formation of mesoderm in *Xenopus* embryos is one of the earliest and most intensely studied inductive events in developmental biology (reviewed in Slack, 1991). The finding that *Wnt* genes can contribute to the experimental induction of *Xenopus* mesoderm has therefore attracted considerable attention and provided new insights into the biological functions of the genes.

After the polarity of frog embryos has been established by the site of sperm entry and subsequent cortical rotation, cells in the vegetal hemisphere induce mesoderm in the marginal equatorial zone. Mesodermal cells on the ventral side differentiate into muscle and blood cells, while dorsal mesoderm induces neural ectoderm, thereby determining the anterior–posterior axis of the embryo. Transplantation experiments have shown that these dorsal mesodermal cells, which constitute the so-called Spemann organizer, can induce an additional axis at an ectopic site. In the simplest model (Figure 4), dorsal–vegetal cells secrete a substance that induces equatorial mesoderm to form a Spemann organizer, which in turn produces neural inductive factors. UV treatment, which blocks cortical rotation, prevents the formation of dorsal mesoderm and subse-

quent axis formation, and therefore results in a ventralized embryo.

Mesoderm induction can be mimicked in vitro by addition of factors to isolated ectodermal caps; the induced mesoderm can then itself induce the appearance of neural structures, instead of epidermis, in the ectodermal pieces. Alternatively, the injection of appropriate mRNAs into individual blastomeres can provoke formation of an additional axis or rescue UV-treated embryos. Through such experiments, several growth factors from the FGF and the TGF β family, activin in particular, have been implicated as mesoderm-inducing growth factors (Kimelman and Kirschner, 1987; Smith et al., 1990; Thomson et al., 1990; Slack et al., 1989). But activin has been reported to induce only a partial dorsal axis, lacking anterior structures, and activin RNA appears unable to rescue UV-treated embryos (Sokol et al., 1991).

Various members of the *Wnt* gene family, injected as mRNA, have more potent effects. For instance, injection of mouse *Wnt-1* RNA into fertilized eggs causes the primary axis to split anteriorly, producing two-headed embryos (McMahon and Moon, 1989). More impressive, injections of *Wnt* RNA into individual ventral blastomeres in 16- to 32-cell embryos induce a complete secondary axis (Figure 4), including the most anterior structures and sometimes a tertiary axis when an additional blastomere is injected (Sokol et al., 1991). Moreover, UV-treated embryos can be rescued almost completely by *Wnt-1* RNA, just as though functional organizer tissue had been implanted. In line with the supposition that Wnt-1 protein has organizing activity, the *Wnt*-injected cells can sometimes be detected in the induced axis itself when injected into certain cells (Sokol et al., 1991). Injection into other blastomeres, however, can still induce a dorsal axis consisting of uninjected cells, indicating that *Wnt* acts as the dorsal–vegetal inducer or the so-called Nieuwkoop center (Smith and Harland, 1991; Figure 4).

Table 4. *Drosophila* Segment Polarity Genes

Phenotype	Genes	Structure of Protein/Homology to:	Interaction with <i>wg</i>	Maternal Effect?	Autonomous?
<i>naked</i> -like	<i>naked</i>	Not determined	Phenotype of <i>naked</i> and <i>zeste-white-3</i> similar to HS- <i>wg</i> ; inhibit expression of <i>en</i>	?	?
	<i>zeste-white-3</i>	Serine/threonine kinase		Yes	Yes
<i>patched</i> -like	<i>patched</i>	Multiple transmembrane	Inhibits <i>wg</i> expression	No	No
	<i>costal-2</i>	Not determined	In <i>patched</i> pathway	Yes	No
<i>wg</i> -like	<i>wg</i>	Secreted factor		No	No
	<i>arm</i>	β -Catenin/plakoglobin	Required for normal function of <i>wg</i> ; protein localization controlled by <i>wg</i>	Yes	Yes
	<i>porcupine</i>	Not determined	Required for normal function of <i>wg</i>	Yes	No
	<i>dsh</i>	Sequence known, novel structure	Required for normal function of <i>wg</i>	Yes	Yes
	<i>fused</i>	Serine/threonine kinase	In <i>patched</i> pathway	Yes	?
	<i>gooseberry</i>	Homeobox; paired box	Maintains late <i>wg</i> expression	No	No
	<i>hedgehog</i>	Not determined	May relieve <i>patched</i> inhibition of <i>wg</i>	No	No
	<i>Ci/D/Cell</i>	Zinc finger		No	No
	<i>smooth</i>	Not determined		?	?
<i>en</i> -like	<i>en</i>	Homeobox	Expression transiently dependent on <i>wg</i> ; and vice versa	No	Yes

Of the several other members of the *Wnt* gene family tested in these biological assays, *Xenopus Wnt-1* and mouse *Wnt-3A* (R. Moon, personal communication), *wg* (Chakrabarti et al., 1992), and the *Xenopus Wnt-8* RNA can induce a dorsal axis (Table 3). In fact, an *XWnt-8* clone has also been isolated by screening cDNAs from a *Xenopus* expression library for competence to rescue UV-treated embryos (Smith and Harland, 1991). The fact that five *Wnt* genes give the same phenotype suggests that their products are all capable of acting on similar endogenous receptors. *Xenopus Wnt-5A* RNA has different effects: it induces head defects and a posterior duplication of the axis after injection of early embryos (R. Moon, personal communication). This may mean that *Wnt-5A* protein acts through a different *Wnt* receptor.

While these results strongly suggest that *Wnt* genes are involved in the induction of dorsal mesoderm and the subsequent establishment of the body axis, it is not clear how they do so. None of the several *Xenopus Wnt* genes tested is naturally expressed during the determination of the organizer. Some show patterns similar to those found in the mouse: *XWnt-1*, *XWnt-3A*, and *XWnt-4* RNAs, for example, are present in the developing CNS at the neurula stage, and *XWnt-8* RNA is found in the marginal zone of the mid-blastula, but not at the right place (dorsal-vegetal cells) or during the right time to act as an endogenous inducer (Christian et al., 1991a, 1991b; Smith and Harland, 1991; Table 3). *XWnt-5A* is the only one to be provided maternally, but it does not induce axes. Possibly an as yet unidentified member of the *Xenopus Wnt* family is the real mesoderm inducer, acting through a receptor that can cross-react with the products of injected *Wnt* RNA. Alternatively, the biochemical effects of *Wnt* protein might intersect with the signal transduction pathway normally stimulated by an endogenous but different type of inducer.

The ultimate specificity of *Wnt* action as a mesoderm modifier may depend on cooperation with other secreted factors, in the way that some differentiation factors achieve specificity through combinatorial action (Sporn and Rob-

erts, 1988). When animal caps isolated from *Xenopus* embryos are treated in culture with bFGF, they produce mesoderm of ventral character; but a similar experiment with animal caps from embryos injected with *XWnt-8* results in mesoderm formation with a distinctly more dorsal nature (Christian et al., 1992). Those explants are also able to produce neural structures not seen after bFGF treatment alone. Since bFGF is present at the right time and place to act as a natural mesoderm inducer (Kimelman et al., 1988), but cannot induce dorsal mesoderm on its own, it has been proposed that *Wnt* modifies the response to bFGF. This observation may be relevant as well to the previously mentioned cooperation between the FGF-related *int-2* gene and *Wnt-1* in virus-induced mammary tumors and in transgenic mice.

Part of the mechanism of action of the *Wnt* genes in *Xenopus* embryogenesis may involve the regulation of gap junctional activity and hence cellular communication. Injection of *Wnt* RNA opens gap junctions at the ventral side of the embryos, where junctions are normally closed (Olson et al., 1991; Olson and Moon, 1992). There is a good correlation between the effects on the anterior dorsal axis and the ventral gap junctions: *Wnt* genes that induce double-headed embryos open junctions, whereas the double tail-inducing *XWnt-5A* fails to do so.

The *Drosophila wg/Wnt* Gene Family

wg, the *Drosophila Wnt-1* ortholog (Rijsewijk et al., 1987a), plays a key role in the establishment of segment polarity in fly embryos. Because of the extensive genetic analysis of *Drosophila* segmentation, insects provide a system particularly amenable to deciphering the machinery of *Wnt* action.

During the first few hours of development, the *Drosophila* embryo is rapidly organized into compartments called parasegments. This process is under the control of a hierarchy of segmentation genes, most of which encode transcription factors (Ingham, 1988). After cellularization, the pattern within each parasegmental unit is further refined

through a diverse group of genes called segment polarity genes. Mutations in those genes lead to deletions and mirror image duplications of the cuticle, usually scored as alterations in the denticle pattern (Nüsslein-Volhard and Wieschaus, 1980). In genetic screens approaching saturation for zygotically required genes, a dozen or so segment polarity genes have been identified and classified according to the region of the cuticle deleted in mutants (Nüsslein-Volhard and Wieschaus, 1980; reviewed in Klingensmith and Perrimon, 1991; Hooper and Scott, 1992 [Table 4]).

Embryonic lethal alleles of *wg* have a very strong segment polarity phenotype and all of the known null mutations map within the protein-encoding domain of *wg* (M. van den Heuvel and R. N., unpublished data). In the absence of the gene, the naked zone of the cuticle is replaced by a continuous lawn of denticles, and parasegmental and segmental boundaries are absent. *wg* null mutations also cause subtle defects in the CNS: the respecification of one neuron present in each segment (Patel et al., 1989). The gene is also required for later development. Several alleles, all of which are rearranged outside the coding domain, are embryonic viable. For example, the first identified allele, *wg*¹, derails the outgrowth of several imaginal discs, in particular the wing disc (Sharma, 1973; Babu, 1977; Baker, 1988), and gave the gene its name.

Early on, it was conjectured that *wg* could be a signaling molecule. Clonal analysis of *wg* mutant cells suggested that the phenotype is nonautonomous in mosaics, i.e., mutant cells can be rescued by surrounding wild-type cells (Morata and Lawrence, 1977; Baker, 1988). The broad zone of deleted cuticle in *wg* mutants relative to the *wg*-expressing area also indicated that *wg* is required for the differentiation of adjacent cells (Baker, 1987). Secretion of the *wg* gene product has subsequently been observed directly (van den Heuvel et al., 1989; González et al., 1991), consistent with its predicted sequence (Rijsewijk et al., 1987a; Cabrera et al., 1987; Udvögyi et al., 1988) and the genetic behavior of *wg*.

In the trunk of the early embryo, *wg* is expressed in 1–2 rows of cells, just anterior to the parasegment boundary (Baker, 1987). This pattern of expression is initiated by earlier acting segmentation genes (such as the pair-rule genes *fushi tarazu* and *even-skipped* [Ingham et al., 1988]), but later becomes dependent on the other segment polarity genes, which interact with each other in a complicated network. Remarkably, the staining patterns for *wg* RNA and protein in whole mount embryos are nearly identical. The protein adheres to the surface of producing cells or associates with the extracellular matrix (van den Heuvel et al., 1989), behaving similarly to its mammalian counterpart and also to a leech *Wnt* gene product (Kostriken and Weisblat, 1992) and to *wg* protein made in *Xenopus* oocytes injected with *wg* RNA (Chakrabarti et al., 1992). But upon close inspection in the electron microscope, some *wg* protein can be detected, presumably endocytosed, in cells adjacent to those that make the RNA (van den Heuvel et al., 1989; González et al., 1991), indicating cell-to-cell spread.




Figure 5. Simplified Scheme of Interactions between the Products of *wg* and Other Segment Polarity Genes

wg positively regulates expression of the *en* (*En*) gene in adjacent cells. The product of *porcupine* (*Porc*), a nonautonomous segment polarity gene, is required for *wg* function. The *wg* receptor is unknown. *armadillo* (*Arm*) and *dishevelled* (*Dsh*) are required in the pathway between *wg* and *en*. The arm protein is homologous to β -catenin and therefore likely present in a complex with cytoskeletal and transmembrane proteins. Transcription of *wg* is negatively regulated by the patched (*Ptc*) protein, a multiple transmembrane protein, through the actions of *fused* (*Fu*), a serine/threonine kinase, and *costal-2* (*Co-2*). Inhibition of patched is relieved by *hedgehog* (*Hh*) activity. Not shown here are many additional regulatory interactions between segment polarity genes.

The *wg* Signal Transduction Pathway

wg influences the expression of many other genes in the *Drosophila* embryo, including *Distal-less*, a homeobox-encoding gene essential for proximal-distal patterning in the limbs (Cohen, 1990), and *labial*, a homeotic gene expressed in the midgut cells adjacent to *wg*-producing visceral mesoderm (Immerglück et al., 1990). The best-known example of such regulatory interactions is the effect of *wg* upon *en* in the epidermis of extended germband embryos (Figure 5). *en* encodes a homeodomain transcription factor that is expressed within the posterior part of each segment, in a row of cells just posterior to those expressing *wg*. Levels of *en* RNA and protein decline prematurely in *wg* mutants (DiNardo et al., 1988; Martínez-Arias et al., 1988; Bejsovec and Martínez-Arias, 1991; Heemskerk et al., 1991); conversely, *en* is necessary for maintenance of *wg* expression. The intercellular signal mediating the control of *wg* by *en* has been postulated to be the product of the *hedgehog* gene, which may interact with the cell surface protein encoded by *patched* (Ingham et al., 1991). Genetic tests indicate that *hedgehog* activates *wg* by preventing the patched cell surface protein from inhibiting expression of *wg* (Figure 5).

wg also controls the localization of the *armadillo* (*arm*) gene product in both an autocrine and a paracrine fashion. *arm* is a segment polarity gene in the *wg* class (Table 4), and it encodes a protein highly similar to the vertebrate proteins plakoglobin and β -catenin, found associated with cadherins in desmosomes and adherens junctions (Peifer and Wieschaus, 1990; McCrea et al., 1991). *arm* RNA and protein are present in all embryonic cells (Riggelman et al., 1989), and the protein colocalizes at the inner cell surface with F-actin, like the junctional proteins to which it is related. In wild type but not in *wg* mutant embryos, the arm protein seems to accumulate to especially high levels in cells making *wg* RNA and in immediately adjacent cells (Riggelman et al., 1990).

When *wg* is expressed ectopically under the control of a heat shock-inducible promoter, the expression of many *wg*-dependent genes changes. In particular, the areas of cells expressing *en* and *Distal-less* broaden, and arm protein is detected at elevated levels in all cells in the embryo (J. Noordermeer, R. N., and P. Lawrence, unpublished data). This implies that *wg* receptors are widespread.

In spite of the extensive analysis of *wg* and other segmentation genes, one of the most sought-after components in this signaling system has not been identified: the *wg* receptor. *arm* and *dishevelled* (*dsh*) both fulfil some criteria for a Wg receptor gene: mutations behave in a cell-autonomous fashion (Table 4; J. Klingensmith and N. Perrimon, personal communication); *arm* and *dsh* deficiencies appear very similar, if not identical, to *wg* deficiencies; and, being maternally acting genes, they are probably expressed ubiquitously. Moreover, in embryos mutant for *arm* and *dsh*, expression of *en* decays in the same pattern as in *wg* mutants, further indicating that their gene products may indeed be in the same pathway (M. van den Heuvel and R. N., unpublished data). Neither *arm* nor *dsh* protein, however, exhibits the hallmarks of cell surface receptor molecules, such as extracellular domains or catalytic functions (J. Klingensmith and N. Perrimon, personal communication). These proteins might be present in a receptor complex, but most likely do not make direct contact with the *wg* ligand.

It is possible that genetic screens have failed to identify the *wg* receptor because of functional redundancy (the presence of multiple receptor genes) or because of pleiotropic effects through interactions with multiple *Wnt* genes. If pleiotropism is responsible, it could be due to the presence of at least two *wg*-related genes *DWnt-2* and *DWnt-3* (Russell et al., 1992; A. McMahon and A. M. C. Brown, personal communication). Transcripts of both genes are detected in early embryos and in limb primordia. *DWnt-2* is expressed in a predominantly segmented pattern and in the presumptive gonads, whereas *DWnt-3* RNA is found in mesodermal and neurogenic regions. If the *DWnt-2* and *DWnt-3* proteins interact with the same receptor as *wg* protein, receptor mutants would probably not resemble *wg* mutants and the receptor gene might not be classified as a segment polarity gene. A confounding factor may also be that the mutational analysis of *Drosophila* segmentation genes has not been exhausted for genes with a maternal contribution (Perrimon et al., 1989).

Common Mechanisms of Wnt Gene Action in Diverse Organisms?

The conservation of Wnt protein sequences during a billion years of evolution (see Figure 1) argues that the proteins retain common functional properties, a notion confirmed by the activities of *Wnt* genes transplanted to distantly related organisms. Thus, *Wnt* genes from flies and mice can induce axis duplication in frog embryos, and *wg* can transform mouse epithelial cells.

Such interchangeability of *Wnt* genes is dramatic, but is it instructive? One potentially useful strategy is to ask whether genes that are known to interact in one organism

exhibit homologous relationships in others. For example, the sites of expression of two *en* genes in the mouse overlap those of *Wnt-1* in the early embryonic midbrain (Davis and Joyner, 1988; McMahon et al., 1992), and there is evidence from both mice and flies for the notion that *Wnt* genes are required to maintain, but not to initiate, expression of *en* genes. Unlike the situation in the *Drosophila* embryo, however, where adjacent rows of cells express either *wg* or *en*, some cells in the developing mouse brain must express both genes. Furthermore, although *en* RNA-positive cells are virtually absent in the developing brain of *Wnt-1*-deficient mice, this may be due to the loss of relevant cells rather than to the loss of a positive regulator of *en*. Thus, it remains uncertain whether *en* and *Wnt* genes cross-regulate each other in vertebrate species as they do in the fly embryo.

The pursuit of *Wnt* genes in many organisms offers the hope of perceiving an important universal effect of Wnt proteins in some experimentally favorable setting. The induction of gap junctions by *Wnt* RNA in ventral cells of *Xenopus* blastomeres has raised the possibility that a general function of Wnt proteins might be to promote cell-cell communication, perhaps by stimulating adhesion of cells as a prelude to formation of gap junctions. Such ideas have been nurtured by the evidence that *Drosophila wg* regulates arm protein, which is, in turn, closely related to proteins present in adherens junctions of vertebrate cells. Since the arm homologs, plakoglobin and β -catenin, associate with transmembrane proteins, such as cadherins, it is tempting to consider the possibility that such transmembrane proteins might be Wnt receptors. In this scheme, Wnt proteins would promote cell-cell adhesion by altering cadherin-like receptors, redistributing arm and its relatives, and ultimately forming gap junctions.

There are, however, problems with this hypothesis. Intuitively, abundant and widespread proteins, such as cadherins, might seem poor candidates for Wnt receptors, given the paucity of cultured cell lines that respond to *Wnt* genes. It is also unlikely that *Wnt*'s role in carcinogenesis would be mediated in this way since transformed mammalian cells tend to be deficient in gap junctions, not overendowed with them (e.g., Eghbali et al., 1991). Furthermore, it is not known whether redistribution of arm protein actually accompanies changes in cell adhesion in the fly embryo, as predicted by the model. Finally, gap junctional communication in *Drosophila* imaginal discs does not coincide with sites of *wg* action (Fraser and Bryant, 1985).

Prospects

Wnt genes are now recognized as mediators of cell-cell signaling events that are important during pattern formation and experimental carcinogenesis, but our understanding of these genes is still in its early stages. The full genetic repertoire has probably not been isolated from any organism, the patterns of expression are incompletely defined for any single gene except *wg*, and the consequences of null mutations are known only for one *Wnt* gene in the mouse and one in the fly.

Most importantly, receptors for Wnt proteins, central in-

gradients in the pathways for signaling, are not known. Many of the proposed organismal effects of Wnt proteins in neural development, tumorigenesis, axis formation, and segmentation need to be related to changes observed in individual cells, such as changes in shape, growth, adhesion, and gap junctions. Any hope of understanding the biochemical events that mediate such changes depends upon identifying and characterizing the molecules that interact directly and specifically with Wnt proteins. Unfortunately, the immediate prospects for isolating Wnt receptors are dimmed by the experimentally unfavorable nature of Wnt proteins, by uncertainties about the distribution, abundance, diversity, and biochemical properties of the receptors, and by the possibility that the numerous Wnt proteins might each be acting through a multisubunit complex formed with components of the extracellular matrix. To overcome these difficulties, it may be essential to learn to prepare some biologically active Wnt protein in a soluble, cell-free form. This is likely to require studies of a variety of Wnt family members, generation of more Wnt mutants, and better mapping of domains involved in secretion, heparin binding, and signaling. If Wnt receptors can be identified, permitting analysis of biochemical responses to Wnt proteins under simple conditions in cultured cells, it should be possible to define with precision the signaling mechanisms that allow Wnt proteins to promote cell growth, differentiation, or survival in the many biological settings reviewed here.

Acknowledgments

We thank Margaret Fuller, Cynthia Kenyon, Matthew Scott, Arend Sidow, Marc Tessier-Lavigne, and members of our laboratory groups for useful criticisms; several colleagues named in the text for permission to cite unpublished work; and the National Institutes of Health and the Howard Hughes Medical Institute for research support.

References

- Aaronson, S. A. (1991). Growth factors and cancer. *Science* 254, 1146-1153.
- Adams, J. M., and Cory, S. (1991). Transgenic models of tumor development. *Science* 254, 1161-1167.
- Alama, S., Casalbone, P., Agostini, E., and Tato, F. (1985). Differentiation of PC12 pheochromocytoma cells induced by v-src oncogene. *Nature* 316, 557-559.
- Ali, I. U., Merlo, G., Callahan, R., and Lidereau, R. (1989). The amplification unit on chromosome 11q13 in aggressive primary human breast tumors entails the *bcl-1*, *int-2* and *hst* loci. *Oncogene* 4, 89-92.
- Babu, P. (1977). Early developmental subdivisions of the wing disk in *Drosophila*. *Mol. Gen. Genet.* 151, 289-294.
- Baker, N. E. (1987). Molecular cloning of sequences from *wingless*, a segment polarity gene in *Drosophila*: the spatial distribution of a transcript in embryos. *EMBO J.* 6, 1765-1773.
- Baker, N. E. (1988). Embryonic and imaginal requirements for *wingless*, a segment-polarity gene in *Drosophila*. *Dev. Biol.* 125, 96-108.
- Bejaouev, A., and Martinez-Arias, A. (1991). Roles of *wingless* in patterning the larval epidermis of *Drosophila*. *Development* 113, 471-485.
- Bittner, J. J. (1936). Some possible effects of nursing on the mammary gland tumor incidence in mice. *Science* 84, 162.
- Blasband, A., Schryver, B., and Papkoff, J. (1992). The biochemical properties and transforming potential of human Wnt-2 are similar to Wnt-1. *Oncogene*, in press.
- Bradley, R. S., and Brown, A. M. C. (1990). The proto-oncogene *int-1* encodes a secreted protein associated with the extracellular matrix. *EMBO J.* 9, 1569-1575.
- Brown, A. M. C., Wildin, R. S., Prendergast, T. J., and Varmus, H. E. (1986). A retrovirus vector expressing the putative mammary oncogene *int-1* causes partial transformation of a mammary epithelial cell line. *Cell* 46, 1001-1009.
- Brown, A. M. C., Papkoff, J., Fung, Y. K., Shackelford, G. M., and Varmus, H. E. (1987). Identification of protein products encoded by the proto-oncogene *int-1*. *Mol. Cell. Biol.* 7, 3971-3977.
- Busse, U., Guay, J., and Séguin, C. (1990). Nucleotide sequence of a cDNA encoding Wnt-1 of the Mexican axolotl *Ambystoma mexicanum*. *Nucl. Acids Res.* 18, 7439.
- Cabrera, C. V., Alonso, M. C., Johnston, P., Phillips, R. G., and Lawrence, P. A. (1987). Phenocopies induced with antisense RNA identify the *wingless* gene. *Cell* 50, 659-663.
- Cantley, L. C., Auger, K. R., Carpenter, C., Duckworth, B., Graziani, A., Kapeller, R., and Soltoff, S. (1991). Oncogenes and signal transduction. *Cell* 64, 281-302.
- Chakrabarti, A., Matthews, G., Dale, L., and Colman, A. (1992). Secretory and inductive properties of the *Drosophila* *wingless* protein in *Xenopus* oocytes and embryos. *Development*, in press.
- Christian, J., Gavin, B., McMahon, A., and Moon, R. (1991a). Isolation of cDNAs partially encoding four *Xenopus* *Wnt-1/int-1* related proteins and characterization of their transient expression during embryonic development. *Dev. Biol.* 143, 230-238.
- Christian, J. L., McMahon, J. A., McMahon, A. P., and Moon, R. T. (1991b). *Xwnt-8*, a *Xenopus* *Wnt-1/int-1*-related gene responsive to mesoderm-inducing growth factors, may play a role in ventral mesodermal patterning during embryogenesis. *Development* 111, 1045-1055.
- Christian, J. L., Olsen, D. J., and Moon, R. T. (1992). *Xwnt-8* modifies the character of mesoderm induced by bFGF in isolated *Xenopus* ectoderm. *EMBO J.* 11, 33-41.
- Cohen, J. C., Shank, P. R., Morris, V. L., Cardiff, R., and Varmus, H. E. (1979). Integration of the DNA of mouse mammary tumor virus in virus-infected normal and neoplastic tissue of the mouse. *Cell* 16, 333-345.
- Cohen, S. M. (1990). Specification of limb development in the *Drosophila* embryo by positional cues from segmentation genes. *Nature* 343, 173-177.
- Davis, C. A., and Joyner, A. L. (1988). Expression patterns of the homeobox-containing genes *En-1* and *En-2* and the proto-oncogene *int-1* diverge during mouse development. *Genes Dev.* 2, 1736-1744.
- Dickson, C., and Peters, G. (1987). Potential oncogene product related to growth factors. *Nature* 326, 833.
- Dickson, C., Smith, R., Brookes, S., and Peters, G. (1984). Tumorigenesis by mouse mammary tumor virus: proviral activation of a cellular gene in the common integration region *int-2*. *Cell* 37, 529-536.
- DiNardo, S., Sher, E., Heemskerk-Jongens, J., Kassiss, J. A., and O'Farrell, P. H. (1988). Two-tiered regulation of spatially patterned *engrailed* gene expression during *Drosophila* embryogenesis. *Nature* 332, 604-609.
- Dollé, P., Izpisua-Belmonte, J.-C., Falkenstein, H., Renucci, A., and Duboule, D. (1989). Coordinate expression of the murine *Hox-5* complex homeobox-containing genes during limb pattern formation. *Nature* 342, 767-772.
- Eghbali, B., Kessler, J. A., Reid, L. M., Roy, C., and Spray, D. C. (1991). Involvement of gap junctions in tumorigenesis: transfection of tumor cells with connexin 32 cDNA retards growth *in vivo*. *Proc. Natl. Acad. Sci. USA* 88, 10701-10705.
- Forrester, L. M., Brunkow, M., and Bernstein, A. (1992). Proto-oncogenes in mammalian development. *Curr. Opin. Genet. Dev.* 2, 38-44.
- Fraser, S. E., and Bryant, P. J. (1985). Patterns of dye coupling in the imaginal wing disk of *Drosophila melanogaster*. *Nature* 317, 533-536.
- Fung, Y. K., Shackelford, G. M., Brown, A. M., Sanders, G. S., and Varmus, H. E. (1985). Nucleotide sequence and expression *in vitro* of cDNA derived from mRNA of *int-1*, a provirally activated mouse mammary oncogene. *Mol. Cell. Biol.* 5, 3337-3344 (Erratum 6, 1362 [1986]).

- Gallahan, D., and Callahan, R. (1987). Mammary tumorigenesis in feral mice: identification of a new *int* locus in mouse mammary tumor virus (Czech II)-induced mammary tumors. *J. Virol.* 61, 66-74.
- Gavin, B. J., and McMahon, A. P. (1992). Differential regulation of the *Wnt*-gene family during pregnancy and lactation suggests a role in post natal development of the mammary gland. *Mol. Cell. Biol.* 12, 2418-2423.
- Gavin, B. J., McMahon, J. A., and McMahon, A. P. (1990). Expression of multiple novel *Wnt-1/int-1*-related genes during fetal and adult mouse development. *Genes Dev.* 4, 2319-2332.
- González, F., Swales, L., Bejsovec, A., Skaer, H., and Arias, A. M. (1991). Secretion and movement of *wingless* protein in the epidermis of the *Drosophila* embryo. *Mech. Dev.* 35, 43-54.
- Hallonet, M. E. R., Teillet, M.-A., and Le Douarin, N. M. (1990). A new approach to the development of the cerebellum provided by the quail-chick marker system. *Development* 108, 19-31.
- Hayward, W. S., Neel, B. G., and Astrin, S. M. (1981). Activation of a cellular oncogene by promoter insertion in ALV-induced lymphoid leukemia. *Nature* 290, 475-480.
- Heemskerk, J., DiNardo, S., Kostriken, R., and O'Farrell, P. H. (1991). Multiple modes of *engrailed* regulation in the progression towards cell fate determination. *Nature* 352, 404-410.
- Hoffmann, F. M., Sternberg, P. W., and Herskowitz, I. (1992). Learning about cancer genes through invertebrate genetics. *Curr. Opin. Genet. Dev.* 2, 45-52.
- Hooper, J., and Scott, M. (1992). The Molecular Genetic Basis of Positional Information in Insect Segments. In *Results and Problems in Cell Differentiation*, W. Hennig, ed. (Berlin: Springer-Verlag).
- Immerglück, K., Lawrence, P. A., and Bienz, M. (1990). Induction across germ layers in *Drosophila* mediated by a genetic cascade. *Cell* 62, 261-268.
- Ingham, P. W. (1988). The molecular genetics of embryonic pattern formation in *Drosophila*. *Nature* 335, 25-34.
- Ingham, P. W., Baker, N. E., and Martínez-Arias, A. (1988). Regulation of segment polarity genes in the *Drosophila* blastoderm by *fushi tarazu* and *even-skipped*. *Nature* 331, 73-75.
- Ingham, P. W., Taylor, A. M., and Nakano, Y. (1991). Role of the *Drosophila patched* gene in positional signalling. *Nature* 353, 184-187.
- Jakobovits, A., Shackleford, G. M., Varmus, H. E., and Martin, G. R. (1986). Two proto-oncogenes implicated in mammary carcinogenesis, *int-1* and *int-2*, are independently regulated during mouse development. *Proc. Natl. Acad. Sci. USA* 83, 7806-7810.
- Jue, S. F., Bradley, R. S., Rudnicki, J. A., Varmus, H. E., and Brown, A. M. C. (1992). The mouse *Wnt-1* gene can act via a paracrine mechanism in transformation of mammary epithelial cells. *Mol. Cell. Biol.* 12, 321-328.
- Kamb, A., Weir, M., Rudy, B., Varmus, H., and Kenyon, C. (1989). Identification of genes from pattern formation, tyrosine kinase, and potassium channel families by DNA amplification. *Proc. Natl. Acad. Sci. USA* 86, 4372-4376.
- Klefer, P., Peters, G., and Dickson, C. (1991). The *Int-2/Fgf-3* oncogene product is secreted and associates with extracellular matrix: implications for cell transformation. *Mol. Cell. Biol.* 11, 5929-5936.
- Kimelman, D., and Kirschner, M. (1987). Synergistic induction of mesoderm by FGF and TGF- β and the identification of an mRNA coding for FGF in the early *Xenopus* embryo. *Cell* 51, 869-877.
- Kimelman, D., Abraham, J. A., Haaparanta, T., Palisi, T. M., and Kirschner, M. W. (1988). The presence of fibroblast growth factor in the frog egg: its role as a natural mesoderm inducer. *Science* 242, 1053-1056.
- Kitajewski, J., Mason, J. O., and Varmus, H. E. (1992). Interaction of the *Wnt-1* proteins with the binding protein BIP. *Mol. Cell. Biol.* 12, 784-790.
- Klagsbrun, M., and Baird, A. (1991). A dual receptor system is required for basic fibroblast growth factor activity. *Cell* 67, 229-232.
- Klingensmith, J., and Perrimon, N. (1991). Segment polarity genes and intercellular communication in *Drosophila*. In *Cell Activation: Genetic Approaches* (New York: Raven Press), pp. 251-274.
- Klingensmith, J., Noll, E., and Perrimon, N. (1989). The segment polarity phenotype of *Drosophila* involves differential tendencies toward transformation and cell death. *Dev. Biol.* 134, 130-145.
- Korteweg, R. (1936). On the manner in which the disposition to carcinoma of the mammary gland is inherited in mice. *Genetics* 18, 350-371.
- Kostriken, R., and Weisblat, D. (1992). Expression of a *Wnt* gene in embryonic epithelium of the leech. *Dev. Biol.* 151, 225-241.
- Kwan, H., Pecanka, V., Tsukamoto, A., Parslow, T. G., Guzman, R., Lin, T.-P., Muller, W. J., Lee, F. S., Leder, P., and Varmus, H. E. (1992). Transgenes expressing the *Wnt-1* and *int-2* proto-oncogenes cooperate during mammary carcinogenesis in doubly transgenic mice. *Mol. Cell. Biol.* 12, 147-154.
- Lammie, G. A., Fantl, V., Smith, R., Schuurin, E., Brookes, S., Michalides, R., Dickson, C., Arnold, A., and Peters, G. (1991). D11S287, a putative oncogene on chromosome 11q13, is amplified and expressed in squamous cell and mammary carcinomas and linked to BCL-1. *Oncogene* 6, 439-444.
- Lane, P. W. (1967). *Mouse Newsletter* 36, 40.
- Lewin, B. (1991). Oncogenic conversion by regulatory changes in transcription factors. *Cell* 64, 303-312.
- Lumsden, A., and Wilkinson, D. (1990). Developmental biology. The promise of gene ablation. *Nature* 347, 335-336.
- Martínez-Arias, A., Baker, N. E., and Ingham, P. W. (1988). Role of segment polarity genes in the definition and maintenance of cell states in the *Drosophila* embryo. *Development* 103, 157-170.
- Mason, J. O., Kitajewski, J., and Varmus, H. E. (1992). Mutational analysis of mouse *Wnt-1* identifies two temperature-sensitive alleles and attributes of *Wnt-1* protein essential for transformation of a mammary epithelial cell line. *Mol. Biol. Cell*, in press.
- McCrea, P. D., Turck, C. W., and Gumbiner, B. (1991). A homolog of the *armadillo* protein in *Drosophila* (plakoglobin) associated with E-cadherin. *Science* 254, 1359-1361.
- McMahon, A. P., and Bradley, A. (1990). The *Wnt-1 (int-1)* proto-oncogene is required for development of a large region of the mouse brain. *Cell* 62, 1073-1085.
- McMahon, A. P., and Moon, R. T. (1989). Ectopic expression of the proto-oncogene *int-1* in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* 58, 1075-1084.
- McMahon, A. P., Joyner, A. L., Bradley, A., and McMahon, J. A. (1992). The midbrain-hindbrain phenotype of *Wnt-1^{+/+}Wnt-1^{-/-}* mice results from stepwise deletion of *engrailed*-expressing cells by 9.5 days postcoitum. *Cell* 69, 581-595.
- McMahon, J. A., and McMahon, A. P. (1989). Nucleotide sequence, chromosomal localization and developmental expression of the mouse *int-1*-related gene. *Development* 107, 643-650.
- Mester, J., Wagenaar, E., Sluysers, M., and Nusse, R. (1987). Activation of *int-1* and *int-2* mammary oncogenes in hormone-dependent and -independent mammary tumors of GR mice. *J. Virol.* 61, 1073-1078.
- Molven, A., Njåstad, P., and Fjose, A. (1991). Genomic structure and restricted neural expression of the zebrafish *wnt-1 (int-1)* gene. *EMBO J.* 10, 799-807.
- Morata, G., and Lawrence, P. A. (1977). The development of *wingless*, a homeotic mutation of *Drosophila*. *Dev. Biol.* 58, 227-240.
- Noordermeer, J., Meijlink, F., Verrijzer, P., Rijsewijk, F., and Destree, O. (1989). Isolation of the *Xenopus* homolog of *int-1/wingless* and expression during neurula stages of early development. *Nucl. Acids Res.* 17, 11-18.
- Nusse, R., and Varmus, H. E. (1982). Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. *Cell* 31, 99-109.
- Nusse, R., van Ooyen, A., Cox, D., Fung, Y. K., and Varmus, H. (1984). Mode of proviral activation of a putative mammary oncogene (*int-1*) on mouse chromosome 15. *Nature* 307, 131-136.
- Nusse, R., Theunissen, H., Wagenaar, E., Rijsewijk, F., Gennissen, A., Otte, A., Schuurin, E., and van Ooyen, A. (1990). The *Wnt-1 (int-1)* oncogene promoter and its mechanism of activation by insertion of proviral DNA of the mouse mammary tumor virus. *Mol. Cell. Biol.* 10, 4170-4179.

- Nusse, R., Brown, A., Papkoff, J., Scambler, P., Shackleford, G., McMahon, A., Moon, R., and Varmus, H. (1991). A new nomenclature for *int-1* and related genes: the *Wnt* gene family. *Cell* 64, 231.
- Nüsslein-Volhard, C., and Wieschaus, E. (1980). Mutations affecting segment number and polarity in *Drosophila*. *Nature* 287, 795–801.
- Oliver, G., Wright, C. V. E., Hardwicke, J., and De Robertis, E. M. (1988). A gradient of homeodomain protein in developing forelimbs of *Xenopus* and mouse embryos. *Cell* 55, 1017–1024.
- Olson, D. J., and Moon, R. T. (1992). Distinct effect of ectopic expression of *Wnt-1*, activin B, and bFGF on gap junctional permeability in 32-cell *Xenopus* embryos. *Dev. Biol.* 151, 204–212.
- Olson, D. J., Christian, J. L., and Moon, R. T. (1991). Effect of *Wnt-1* and related proteins on gap junctional communication in *Xenopus* embryos. *Science* 252, 1173–1176.
- Papkoff, J. (1989). Inducible overexpression and secretion of *int-1* protein. *Mol. Cell. Biol.* 9, 3377–3384.
- Papkoff, J., and Schryver, B. (1990). Secreted *int-1* protein is associated with the cell surface. *Mol. Cell. Biol.* 10, 2723–2730.
- Papkoff, J., Brown, A. M. C., and Varmus, H. E. (1987). The *int-1* proto-oncogene products are glycoproteins that appear to enter the secretory pathway. *Mol. Cell. Biol.* 7, 3978–3984.
- Patel, N. H., Schaller, B., Goodman, C. S., and Holmgren, R. (1989). The role of segment polarity genes during *Drosophila* neurogenesis. *Genes Dev.* 3, 890–904.
- Peifer, M., and Wieschaus, E. (1990). The segment polarity gene *armadillo* encodes a functionally modular protein that is the *Drosophila* homolog of human plakoglobin. *Cell* 63, 1167–1178.
- Perkins, A., Kongsuwan, K., Visvader, J., Adams, J., and Cory, S. (1990). Homeobox gene expression plus autocrine growth factor production elicits myeloid leukemia. *Proc. Natl. Acad. Sci. USA* 87, 8398–8402.
- Perrimon, N., Engstrom, L., and Mahowald, A. P. (1989). Zygotic lethals with specific maternal effect phenotypes in *Drosophila melanogaster*. I. Loci on the X chromosome. *Genetics* 121, 333–352.
- Peters, G., Lee, A. E., and Dickson, C. (1986). Concerted activation of two potential proto-oncogenes in carcinomas induced by mouse mammary tumour virus. *Nature* 320, 628–631.
- Peters, G., Brookes, S., Smith, R., Placzek, M., and Dickson, C. (1989). The mouse homolog of the *hst/k-FGF* gene is adjacent to *int-2* and is activated by proviral insertion in some virally induced mammary tumors. *Proc. Natl. Acad. Sci. USA* 86, 5678–5682.
- Rabbitts, T. H. (1991). Translocations, master genes, and differences between the origins of acute and chronic leukemias. *Cell* 67, 641–644.
- Riggleman, B., Wieschaus, E., and Schedl, P. (1989). Molecular analysis of the *armadillo* locus: uniformly distributed transcripts and a protein with novel internal repeats are associated with a *Drosophila* segment polarity gene. *Genes Dev.* 3, 96–113.
- Riggleman, B., Schedl, P., and Wieschaus, E. (1990). Spatial expression of the *Drosophila* segment polarity gene *armadillo* is posttranscriptionally regulated by *wingless*. *Cell* 63, 549–560.
- Rijsewijk, F., Schuermann, M., Wagenaar, E., Parren, P., Weigel, D., and Nusse, R. (1987a). The *Drosophila* homolog of the mouse mammary oncogene *int-1* is identical to the segment polarity gene *wingless*. *Cell* 50, 649–657.
- Rijsewijk, F., van Deemter, L., Wagenaar, E., Sonnenberg, A., and Nusse, R. (1987b). Transfection of the *int-1* mammary oncogene in cuboidal RAC mammary cell line results in morphological transformation and tumorigenicity. *EMBO J.* 6, 127–131.
- Robbins, J., Blondel, B. J., Gallahan, D., and Callahan, R. (1992). Mouse mammary tumor gene *int-3*: a member of the *notch* gene family transforms mammary epithelial cells. *J. Virol.* 66, 2594–2599.
- Roelink, H., and Nusse, R. (1991). Expression of two members of the *Wnt* family during mouse development—restricted temporal and spatial patterns in the developing neural tube. *Genes Dev.* 5, 381–388.
- Roelink, H., Wagenaar, E., Lopes da Silva, S., and Nusse, R. (1990). *Wnt-3*, a gene activated by proviral insertion in mouse mammary tumors, is homologous to *int-1/Wnt-1* and normally expressed in mouse embryos and adult brain. *Proc. Natl. Acad. Sci. USA* 87, 4519–4523.
- Roelink, H., Wagenaar, E., and Nusse, R. (1992). Amplification and proviral activation of several *Wnt* genes during progression and clonal variation of mouse mammary tumors. *Oncogene* 7, 487–492.
- Ruangvoravat, C. P., and Lo, C. W. (1992). Restrictions in gap junctional communication in the *Drosophila* larval epidermis. *Dev. Dynamics* 193, 70–82.
- Russell, J., Gennissen, A., and Nusse, R. (1992). Isolation and expression of two novel *Wnt/wingless* gene homologs in *Drosophila*. *Development*, in press.
- St. Arnaud, R., Craig, J., McBurney, M. W., and Papkoff, J. (1989). The *int-1* proto-oncogene is transcriptionally activated during neuroectodermal differentiation of P19 mouse embryonal carcinoma cells. *Oncogene* 4, 1077–1080.
- Schuuring, E., van Deemter, L., Roelink, H., and Nusse, R. (1989). Transient expression of the proto-oncogene *int-1* during differentiation of P19 embryonal carcinoma cells. *Mol. Cell. Biol.* 9, 1357–1361.
- Shackleford, G. M., and Varmus, H. E. (1987). Expression of the proto-oncogene *int-1* is restricted to postmeiotic male germ cells and the neural tube of mid-gestational embryos. *Cell* 50, 89–95.
- Sharma, R. P. (1973). *wingless*, a new mutant in *D. melanogaster*. *Dros. Inf. Serv.* 50, 134.
- Sidow, A. (1992). Diversification of the *Wnt* gene family on the ancestral lineage of vertebrates. *Proc. Natl. Acad. Sci. USA*, in press.
- Slack, J. (1991). *From Egg to Embryo* (Cambridge: Cambridge University Press).
- Slack, J. M. W., Darlington, B. G., Gillespie, L. L., Godsave, S. F., Isaacs, H. V., and Paterno, G. D. (1989). The role of fibroblast growth factor in early *Xenopus* development. *Development (Suppl.)* 707, 141–148.
- Smith, J. C., Price, B. M. J., van Nimmen, K., and Huylebroeck, D. (1990). Identification of a potent *Xenopus* mesoderm-inducing factor as a homolog of activin A. *Nature* 345, 729–731.
- Smith, W. C., and Harland, R. M. (1991). Injected *Xwnt-8* RNA acts early in *Xenopus* embryos to promote formation of a vegetal dorsalizing center. *Cell* 67, 753–765.
- Sokol, S., Christian, J. L., Moon, R. T., and Melton, D. A. (1991). Injected *Wnt* RNA induces a complete body axis in *Xenopus* embryos. *Cell* 67, 741–752.
- Sonnenberg, A., van Balen, P., Hilgers, J., Schuuring, E., and Nusse, R. (1987). Oncogene expression during progression of mouse mammary tumor cells; activity of a proviral enhancer and the resulting expression of *int-2* is influenced by the state of differentiation. *EMBO J.* 6, 121–125.
- Sporn, M. B., and Roberts, A. B. (1988). Peptide growth factors are multifunctional. *Nature* 332, 217–219.
- Stewart, T. A., Pattengale, P. K., and Leder, P. (1984). Spontaneous mammary adenocarcinomas in transgenic mice that carry and express *MTV/myc* fusion genes. *Cell* 38, 627–637.
- Taira, M., Yoshida, T., Miyagawa, K., Sakamoto, H., Terada, M., and Sugimura, T. (1987). cDNA sequence of human transforming gene *hst* and identification of the coding sequence required for transforming activity. *Proc. Natl. Acad. Sci. USA* 84, 2980–2984.
- Thomas, K. R., and Capecchi, M. R. (1990). Targeted disruption of the murine *int-1* proto-oncogene resulting in severe abnormalities in midbrain and cerebellar development. *Nature* 346, 847–850.
- Thomas, K. R., Musci, T. S., Neumann, P. E., and Capecchi, M. R. (1991). *Swaying* is a mutant allele of the proto-oncogene *Wnt-1*. *Cell* 67, 969–976.
- Thomsen, G., Woolf, T., Whitman, M., Sokol, S., Vaughan, J., Vale, W., and Melton, D. A. (1990). Activins are expressed early in *Xenopus* embryogenesis and can induce axial mesoderm and anterior structures. *Cell* 63, 485–493.
- Tsukamoto, A. S., Grosschedl, R., Guzman, R. C., Parslow, T., and Varmus, H. E. (1988). Expression of the *int-1* gene in transgenic mice is associated with mammary gland hyperplasia and adenocarcinomas in male and female mice. *Cell* 55, 619–625.
- Uzvolgyi, E., Kiss, I., Pitt, A., Arsenian, S., Ingvarsson, S., Udvardy, A., Hamada, M., Klein, G., and Sümegi, J. (1988). *Drosophila* homolog

of the murine *Int-1* protooncogene. *Proc. Natl. Acad. Sci. USA* 85, 3034–3038.

van den Heuvel, M., Nusse, R., Johnston, P., and Lawrence, P. A. (1989). Distribution of the *wingless* gene product in *Drosophila* embryos: a protein involved in cell–cell communication. *Cell* 59, 739–749.

Van de Vijver, M. J., and Nusse, R. (1991). The molecular biology of breast cancer. *Biochim. Biophys. Acta* 1072, 33–50.

Van de Vijver, M. J., Petersen, J., Mool, W., Lomans, J., Devilee, P., Cornelisse, C., Bos, J. L., and Nusse, R. (1989). Oncogene activations in human breast cancer. *Cancer Cells* 7, 385–391.

van Lohulzen, M., and Berns, A. (1990). Tumorigenesis by slow-transforming retroviruses—an update. *Biochim. Biophys. Acta* 1032, 213–235.

van Ooyen, A., and Nusse, R. (1984). Structure and nucleotide sequence of the putative mammary oncogene *int-1*; proviral insertions leave the protein-encoding domain intact. *Cell* 39, 233–240.

van Ooyen, A., Kwee, V., and Nusse, R. (1985). The nucleotide sequence of the human *int-1* mammary oncogene; evolutionary conservation of coding and non-coding sequences. *EMBO J.* 4, 2905–2909.

Van't Veer, L. J., Geurts van Kessel, A., Van Heerikhuizen, H., van Ooyen, A., and Nusse, R. (1984). Molecular cloning and chromosomal assignment of the human homolog of *int-1*, a mouse gene implicated in mammary tumorigenesis. *Mol. Cell. Biol.* 4, 2532–2534.

Wainwright, B. J., Scambler, P. J., Stanier, P., Watson, E. K., Bell, G., Wicking, C., Estivill, X., Courtney, M., Boue, A., Pedersen, P. S., Williamson, R., and Farrall, M. (1988). Isolation of a human gene with protein sequence similarity to human and murine *int-1* and the *Drosophila* segment polarity mutant *wingless*. *EMBO J.* 7, 1743–1748.

Wilkinson, D. G., Bailes, J. A., and McMahon, A. P. (1987). Expression of the proto-oncogene *int-1* is restricted to specific neural cells in the developing mouse embryo. *Cell* 50, 79–88.